arXiv:hep-ph/0312384v1 31 Dec 2003

# Quick View on Jet Tomography at RHIC Energies

G. G. Barnaföldi[a] *

[a]RMKI KFKI, P.O. Box 49, Budapest 1525, Hungary;
ITOL, Eötvös University, Pázmány P. 1/A, Budapest 1117, Hungary

A strong suppression of $\pi$ production in $AuAu$ collision was discovered at RHIC energies in the high-$p_T$ region. The lack of this suppression in $dAu$ collision proved, this effect can be connected to final state effects, namely medium induced jet energy loss caused by dense color matter. Here we display our quantitative results on jet quenching in $AuAu$ collision using our pQCD-based parton model calculations. In parallel we present results on $dAu$ collision.

## 1. Short introduction

Recent high-$p_T$ RHIC experimantal results on pion production in central $AuAu$ collision at mid-rapidity at $\sqrt{s} = 130$ and 200 AGeV have shown a strong suppression compared to binary scaled $pp$ data [ 1, 2]. Analysing this effect in different impact parameter ranges, the suppression vanishes with increasing centrality. In $dAu$ the lack of the suppression [ 1] was found also. A quantitative analysis of the suppression pattern, the so called *"jet-tomography"* [ 3], yields information about the physical aspects of the produced dense matter and the impact parameter dependence of the formation of deconfined matter in $AuAu$ collisions. Latest data on $dAu$ collision confirmed that initial state effects can not be responsible for suppression in $AuAu$ collision. Thus medium induced non-Abelian jet-energy loss in the final state becomes a strong candidate to explain the missing pion yield. Jet energy loss can be calculated in a perturbative quantum chromodynamics (pQCD) frame [ 4]. Our aim is to calculate $\pi$ production in nucleus-nucleus $AA$ collision in a pQCD improved parton model, taking into account a phenomenological intrinsic transverse momentum distribution (intrinsic-$k_T$) for the colliding partons, which is necessary to get better agreement between data and calculations in $pp$ collisions [ 5, 6]. For $AA$ collision we included initial state effects into our calculations: nuclear multiple scattering, saturation in the number of semihard collisions, and shadowing effect inside nucleus. We used Glauber-like collision geometry.

Here we investigate jet energy loss. As high-energy quark and gluon jets are travelling throught dense colored matter, they can lost their energy described by induced gluon

---

*Thanks the EPS Scholarship and OTKA U047552 for support of my participation, and for my co-authors, P. Lévai, G. Papp, G. Fai., for discussions, and advices. This work was supported in part by Hungarian grants T034842, T043455, T043514, U.S. DOE grant: DE-FG02-86ER40251, and NSF grant: INT0000211. Supercomputer time provided by BCPL in Norway and the EC – Access to Research Infrastructure action of the Improving Human Potential Programme is gratefully acknowledged.

radiation in thin non-Abelian matter. The GLV-description [ 4] of energy loss can be included into the pQCD improved parton model as a strong final state effect. Comparing theory to data we can extract the opacity values ($\bar{n} = L/\lambda$) of the produced colored matter at different centralities.

## 2. Numerical results vs. experimental data

Applying the above sketched model for reproducing $\pi$ yields in $AuAu$, and $dAu$ collisions we can accomplish a tomographic analysis of the produced dense and colored matter in reactions. In Fig. 1 we display the nuclear modification factor: $R_{AA'}(p_T) = \frac{E_\pi \mathrm{d}\sigma_\pi^{AA'}/\mathrm{d}^3p_T}{N_{bin}\ E_\pi \mathrm{d}\sigma_\pi^{pp}/\mathrm{d}^3p_T}$, as a function of $p_T$ for pion production from central to peripheral $AuAu$ (*lower curves*) and in $dAu$ (*the upper curve*). In the most central $AuAu$ collision the measured suppression is reproduced with opacity $L/\lambda = 3.5$. As Fig. 1 displays, in $AuAu$ collisions the suppression of the high-energy pions goes away with decreasing centrality, which is followed by the descreasing values of the opacity. In $dAu$ collision we do not need to include jet quenching to reproduce pion production, this final state effect does not appear there, only multiscattering and shadowing.



Figure 1. $R_{AA}$ for pions from $AuAu$ and $dAu$ collisions, data are taken from Refs. [ 1, 2], and calculation for $dAu$ from Ref. [ 7].

## REFERENCES

1. G. David *et al.* (PHENIX Coll.), Nucl. Phys. A698 (2002) 227; D. d'Enterria *et al.* (PHENIX Coll.), Nucl. Phys. A715 (2003) 749; S.S. Adler *et al.* (PHENIX Coll.), Phys. Rev. Lett. 91 (2003) 072301, *ibid.* 072303; `nucl-ex/0308006`.
2. J.C. Dunlop *et al.* (STAR Coll.), Nucl. Phys. A698 (2002) 515; J.L. Klay *et al.* (STAR Coll.), Nucl. Phys. A715 (2003) 733; J. Adams *et al.* (STAR Coll.), Phys. Rev. Lett. 91 (2003) 172302, *ibid.* 072304; `nucl-ex/0308023`.
3. E. Wang, X.N. Wang, Phys. Rev. Lett. 89 (2002) 162301; M. Gyulassy, P. Lévai, I. Vitev, Phys. Lett. B538 (2002) 282.
4. M. Gyulassy, P. Lévai and I. Vitev, Phys. Rev. Lett. 85 (2000) 5535; Nucl. Phys. B571 (2000) 197, *ibid.* B594 (2001) 371.
5. G. Papp et al. `hep-ph/0212249`; Y. Zhang *et al.*, Phys. Rev. C65 (2002) 034903.
6. G.G. Barnaföldi *et al.*, APH NS Heavy Ion Phys. 18 (2003) 79, `nucl-th/0206006`.
7. P. Lévai. *et al.*, `nucl-th/0306019`.